

Review

CPLANE Complex and Ciliopathies

Jesús Eduardo Martín-Salazar¹ and Diana Valverde^{1,2,*} ¹ CINBIO, Biomedical Research Centre, University of Vigo, 36310 Vigo, Spain; edu19ms@gmail.com² Galicia Sur Health Research Institute (IIS-GS), 36310 Vigo, Spain

* Correspondence: dianaval@uvigo.es

Abstract: Primary cilia are non-motile organelles associated with the cell cycle, which can be found in most vertebrate cell types. Cilia formation occurs through a process called ciliogenesis, which involves several mechanisms including planar cell polarity (PCP) and the Hedgehog (Hh) signaling pathway. Some gene complexes, such as BBSome or CPLANE (ciliogenesis and planar polarity effector), have been linked to ciliogenesis. CPLANE complex is composed of *INTU*, *FUZ* and *WDPCP*, which bind to *JBTS17* and *RSG1* for cilia formation. Defects in these genes have been linked to a malfunction of intraflagellar transport and defects in the planar cell polarity, as well as defective activation of the Hedgehog signalling pathway. These faults lead to defective cilium formation, resulting in ciliopathies, including orofacial–digital syndrome (OFDS) and Bardet–Biedl syndrome (BBS). Considering the close relationship, between the CPLANE complex and cilium formation, it can be expected that defects in the genes that encode subunits of the CPLANE complex may be related to other ciliopathies.

**Citation:** Martín-Salazar, J.E.;Valverde, D. CPLANE Complex and Ciliopathies. *Biomolecules* **2022**, *12*, 847.[https://doi.org/10.3390/](https://doi.org/10.3390/biom12060847)[biom12060847](https://doi.org/10.3390/biom12060847)

Academic Editor: Maliha Zahid

Received: 10 May 2022

Accepted: 16 June 2022

Published: 17 June 2022

Publisher’s Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Correction Statement: This article has been republished with a minor change. The change does not affect the scientific content of the article and further details are available within the backmatter of the website version of this article.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Keywords: cilia; ciliopathies; CPLANE

1. Introduction

The term “ciliopathies” emerged in 2006 [1] to describe a heterogeneous group of rare genetic disorders that share a common aetiology: defects in cilia and centrosome/basal body structure and/or function

Cilia are highly conserved organelles that project from the surface of almost all eukaryotic cells and have important roles primarily in sensory perception and cell motility [2]. Cilia are classified as motile and sensory or primary cilia [3]. Primary cilia are cell-cycle-associated organelles which have been described as small antennae protruding from the cell surface [4] and are present on almost all mammalian cell types, while motile cilia are assembled by certain cell types such as respiratory epithelia, sperm flagella and fallopian tubes [5].

Primary cilia detect and transduce extracellular signals, such as biochemical and mechanical stimuli, to regulate processes including differentiation and proliferation [6].

Since primary cilia were first discovered in chondrocytes in 1967 [7], an increasing number of studies have focused on exploring their functions, mainly their role in endocytosis, infiltration and apoptosis [8]. Initially, primary cilia had been considered unimportant vestigial structures. However, increasing evidence suggests that primary cilia are functional organelles that can not only detect chemical or mechanical signals in the extracellular environment [9], but can also regulate cell mitosis [10] and signal transduction [11].

The primary cilium is composed of nine microtubule doublets (9 + 0 structure), building the ciliary axoneme. This structure is covered by a membrane that is continuous with the plasma membrane, but it has a particular protein and lipid content that is essential for ciliary activity [12,13].

Primary cilia are dynamic organelles that are continually being assembled and re-absorbed in a process tightly coupled to the cell cycle. The assembly process, called ciliogenesis, is made up of several steps. The axoneme elongates from the basal body,

which is the transformed mother centriole with distal and subdistal appendages. Cilia formation is initiated by the apical migration of the mother centriole to become the basal body, followed by extension of the axoneme microtubules, formation of the transition zone and growth of the cilium by the ciliary trafficking machinery such as intraflagellar transport [14,15]. There are different mechanisms and signalling pathways involved in cilia formation, two of the most important being the planar cell polarity (PCP) and the hedgehog signalling pathway (Hh).

Cell polarization refers to the organised establishment of asymmetries within cells. Just as intracellular functions are compartmentalised into organelles, many cellular functions are made more effective by partitioning along an axis of polarisation. Cell polarisation often leads to specific molecular determinants being localised to specific cellular domains, and the coordination of polarity in tissues is essential for the development of specialised forms and functions in multicellular organisms [16].

The Hedgehog (Hh) signalling pathway is an evolutionarily conserved signal transmission pathway from the cell membrane to the nucleus [17]. It plays an important role in normal embryonic development in invertebrates and vertebrates [18].

Some proteins are grouped to function as a whole forming complex, such as BBsome, and have been associated with ciliogenesis and cell polarity, and the hedgehog signalling pathway. The least known is the CPLANE (ciliogenesis and planar polarity effector) complex [19–21]. Defects in the genes encoding components of this complex have been related to failures of the ciliogenesis process and in planar cell polarity, thus associating the CPLANE complex with these processes [22]. However, the exact role of this complex is not well known, and the objective of this review is to collect information about the CPLANE complex and its function.

2. CPLANE Complex Structure

Throughout the combination of proteomic techniques, genetic analysis and in vivo imaging studies of proteins that have been linked to planar cell polarity, a set of Inturned (Inturned Planar Cell Polarity Protein), Fuzzy (Fuzzy Planar Cell Polarity Protein) and Wdpcp (WD Repeat Containing Planar Cell Polarity Effector) proteins, previously related to the ciliogenesis process [21,23], have been identified as a complex called CPLANE (ciliogenesis and planar cell polarity effector) [24]. These three elements are very closely linked to each other. In addition, it has been found that these three proteins interact with others that promote the assembly of the cilium. A clear interaction has been observed with the Jbts17 (Joubert Syndrome 17) protein (also call C5orf42) [24], previously related to Joubert syndrome and Orofacial–digital syndrome [25,26]. The interaction between the atypical Rab-type guanosine triphosphatase (GTPase) Rsg1 and Fuz was already known [20]. However, a recent study also found a link between Intu and Wdpcp with Rsg1 [24], suggesting that Intu favours the recruitment of Rsg1 in the late steps of ciliogenesis initiation [19,27].

The structure of the complex and the different interactions between the elements of the complex were practically unknown until now. A recent study carried out on mice and human cell lines has revealed that the complex adopts a crescent-like architecture, such that Wdpcp and Rsg1 are located at the ends of the crescent and bind to the central Intu-Fuz heterodimer on opposite sides. This creates a linear-type subunit interaction scheme that can be represented as Wdpcp-Intu-Fuz-Rsg1 [28].

Regarding the structure of the different proteins that conform the complex, Wdpcp (85.084 kDa) folds into a seven-bladed β -propeller belonging to the WD40 family, followed by an array of α helices [28]. Both Intu and Fuz have three longin-like domains (LD). LD1 of Intu (105.648 kDa) and Fuz (45.679 kDa) adopts a canonical longin fold, LD2 adopts a longin-like fold and LD3 adopts a lamp longin-like fold [29]. The core structure of Rsg1 (28.534 kDa) follows the typical small GTPase fold characterised by a central six-stranded concave β sheet surrounded by five α helices [28]. Wdpcp mainly contacts Intu with a large interface area of 2609 Å². The main interaction platform is provided by Intu LD2, forming extensive contacts with the α -helical domain of Wdpcp. Intu contacts Fuz with a

large buried surface area of 1898 Å². Intu-Fuz heterodimer formation is mediated by LD1 and LD3 [28].

Nevertheless, it has been shown that Rsg1 interacts only with Fuz [28], and there is no interaction with the other elements of the complex as suspected [19,24]. The Fuz-Rsg1 interface spans an area of 1099 Å² and consists of Fuz LD1 and LD2 engaging the central β sheet of Rsg1 [28].

3. CPLANE and Ciliogenesis

The relationship between the different elements that conform the CPLANE complex and the process of ciliogenesis has been extensively demonstrated.

First, disruption of the *FUZ* gene has been shown to disrupt ciliogenesis, resulting in shorter cilia, in *Xenopus* and mice [20,21,30]. However, in mice, cilia formation in different cell types such as Meckel's cartilage cells, mesenchymal cells of the notochord and limb buds, and other cell types was not completely disrupted in the absence of *FUZ*, suggesting that other PCP effector genes, such as *INTU* [31] or *WDPCP*, may be involved in this process and compensate for the absence of *FUZ* [32].

In contrast, it was observed that the reduction of the amount of Intu protein by morpholino in *Xenopus* leads to the inefficient formation of cilia in the spinal cord and epidermis [21]. Furthermore, in *INTU* null mutant mouse embryos, primary cilia are observed in most ganglion cells. However, many of these cilia are severely atrophied, suggesting that *INTU* is not required for cilia formation per se, but is required for the formation of morphologically normal ganglion cilia [31]. Additionally, in fibroblast cell cultures, it was observed that there is no cilia formation in *INTU* null mutants [31]. All these data suggest that a clear link exists between *INTU* and the process of ciliogenesis.

Less data are available on the relationship between *WDPCP* and ciliogenesis; however, it has been observed that *WDPCP* is required for the recruitment to the ciliary transition zone of Mks1, Sept2 and Nphp1, three proteins required for ciliogenesis [33], and therefore there appears to be a relationship between *WDPCP* and ciliogenesis.

Finally, in a recent study carried out on mice, the effect of *RSG1* on cilia formation has been observed. Although most cells in the mesenchyme of the limb and adjacent to the neural tube are ciliated in the wildtype, fewer than 30% of mesenchymal cells in *RSG1* mutant mice embryos had cilia [19]. However, that 30% have normal cilia lengths, suggesting that the function of *RSG1* is somehow involved in increasing the efficiency of primary cilia initiation. In this study, the recruitment of Rsg1 to the mother centriole was confirmed to depend on its GTPase activity. Intu, which directly interacts with Rsg1 [24], was detected in *RSG1* mutants. Similarly, Ttbk2, a ciliogenesis-initiating protein, was detected in *INTU* mutants and *RSG1* mutants. In this way, it is proposed that Ttbk2 favours the recruitment of Intu, and this of Rsg1, acting Rsg1 in the last steps of the beginning of ciliogenesis [19]. However, as discussed above, a recent study has shown that Rsg1 only interacts with the CPLANE complex via Fuz [28], so the recruitment mechanism of Rsg1 in ciliogenesis remains unknown.

4. CPLANE and Ciliopathies

Ciliopathies comprise a heterogeneous group of genetic disorders caused by an alteration, which may be functional or structural, of the cilia [1,34].

As previously said, primary cilia are present in many tissues and cell types, which explains the wide range of clinical features associated with these disorders. Although these phenotypes can be related to organ-specific diseases, most ciliopathies are pleiotropic syndromes characterised by showing overlapping phenotypes [1,35]. Frequent cilia-related manifestations are (poly)cystic kidney disease, retinal degeneration, situs inversus, cardiac defects, polydactyly, other skeletal abnormalities, and defects of the central and peripheral nervous system, occurring either isolated or as part of syndromes.

There is a close relationship between the CPLANE complex and proper cilia formation and function. Thus, it is to be expected that mutations in the different components of the

complex will lead to a malformation of the cilium and defects in its function, giving rise to ciliopathies. Here, we will try to gather the existing information on the relationship between the genes that make up the complex with the appearance of some ciliopathy.

First, dominant mutations in *FUZ* have been reported to cause isolated neural tube defects [36]. Another study showed that a frameshift deletion mutation in *FUZ* is associated with the development of short rib and polydactyly syndromes (SRPS) [37] (Table 1). The SRPS group includes six distinct autosomal recessive conditions, including four lethal conditions (Saldino-Noonan syndrome or SRPtype I (OMIM 263530), Majewski syndrome or SRP type II (OMIM 263520), Verma–Naumoff syndrome or SRPtype III (OMIM 263510), and Beemer–Langer syndrome or SRPtype IV (OMIM 26986), and two that are compatible with life: *EVC* (OMIM 225500) and Jeune syndrome (ATD, OMIM 611263, OMIM 613091, OMIM 613819, and OMIM 614376). They are characterised by micromelia, short ribs, hypoplastic thorax, polydactyly (pre- and postaxial), and multiple anomalies of major organs [38].

In a recent study, null mutations in the PCP effector gene, *FUZZY*, were observed to cause profound early renal hypoplasia in mice [39].

For *INTU*, null mutant mice are homozygous lethal at mid-gestation and have been associated with the development of severe polydactyly [31]. In humans, a link has been found between mutations in *INTU* and the occurrence of SRPS [24]. Oral–facial–digital (OFD) syndromes are rare genetic disorders characterised by the association of abnormalities of the face (hypertelorism and low-set ears), oral cavity (lingual hamartoma, abnormal frenulum and lobulated tongue) and extremities (brachydactyly and polydactyly). OFD syndromes also comprise a broad range of additional features that initially led to the clinical delineation of 17 OFD subtypes [40]. Mutations in *INTU* are associated with the ciliopathy phenotype in the OFD type VI syndrome spectrum (OMIM 277170) [41]. Additionally, in another study, *INTU* was proposed as a possible cause of OFD type II syndrome (OMIM 252100) [40] and finally, pathogenic variants of *INTU* have been reported in two patients with OFDS XVII (OMIM 617926) [42] (Table 1).

Table 1. Ciliopathies related to CPLANE complex genes.

Gene	Disease	Reference
<i>INTU</i>	Short-Rib Polydactyly Syndrome (SRPS)	[24]
	Nephronophthisis	[24]
	Oro-facial-Digital Syndrome Type 2? (OFD2)	[40]
	Oro-facial-Digital Syndrome Type 17 (OFD17)	[42]
	Oro-facial-Digital Syndrome Type 6 (OFD6)	[41]
<i>FUZ</i>	Short-Rib Polydactyly Syndrome (SRPS)	[37]
<i>WDPCP</i>	Bardet-Biedl Syndrome (BBS)	[23]
	Meckel-Gruber syndrome (MKS)	[33]
<i>JBTS17</i>	Oro-facial-Digital Syndrome Type 6 (OFD6)	[25]
	Joubert Syndrome (JS)	[24]
	Meckel-Gruber syndrome (MKS)	[43]

All associations between genes and different pathologies have been observed in humans.

On the other hand, variants in *WDPCP* have been mainly related to OFDS [40]. It was found that *WDPCP* loss-of-function mutations may be sufficient to cause Bardet–Biedl Syndrome (BBS, OMIM 209900) [23,44]. BBS is a pleiotropic ciliopathy characterised by six main clinical features: retinal dystrophy, truncal obesity, postaxial polydactyly, urogenital anomalies, renal abnormalities and cognitive impairment [45,46]. A wide range of secondary manifestations has been also reported. Mutations in *WDPCP* cause phenotypes similar to those found in individuals affected by Meckel-Gruber syndrome. (MKS, OMIM 249,000) [33] (Table 1). MKS is a lethal developmental syndrome characterised by

posterior fossa abnormalities, bilateral enlarged cystic kidneys, and hepatic developmental defects that include ductal plate malformation associated with hepatic fibrosis and cysts. A common additional feature is postaxial polydactyly, usually affecting both hands and feet; occasional features are described [47].

Within the CPLANE complex, the gene that, until now, had a greater relationship with the appearance of ciliopathies is *JBTS17* (Table 1). To date, a large number of different pathogenic *JBTS17* mutations have been reported (mostly in patients with Joubert syndrome (JS) but a small part in OFDVI patients) [25,48–50]. JS is diagnosed by a pathognomonic malformation of the midbrain–hindbrain junction which results in the brain imaging finding “molar tooth sign” (MTS). Variable involvement of other organs (such as the eye, kidney, liver, and skeleton) is present in two-thirds of individuals with JS and can manifest at different ages and with variable severity [51]. In two more recent studies, new variants in *JBTS17* associated with JS have been identified [52,53]. Additionally, *JBTS17* has been associated with MKS [43].

The Small GTPase Rsg1 is the only component of the CPLANE complex for which no variant associated with the appearance of any ciliopathy has yet been described.

5. How Does the CPLANE Complex Relate to the Signalling Pathways Required for Ciliogenesis?

As mentioned above, the development of ciliopathies is due to failures during cilium formation, and we have seen the relationship between the CPLANE complex and ciliogenesis. More specifically, ciliogenesis depends on processes and signalling pathways such as intraflagellar transport, planar cell polarity or the Hedgehog signalling pathway, showing a link between these processes and the CPLANE complex.

5.1. Intraflagellar Transport

Ciliogenesis and cilia-mediated signalling both require the function of a highly conserved system of intraflagellar transport (IFT). The function of this system is to transport ciliary cargoes by using specific kinesin and dynein motors to move bidirectionally along the microtubule doublets of the ciliary axoneme [13,54]. Several genetic and biochemical studies have shown that the IFT is subdivided into two complexes: IFT-B, which governs the anterograde traffic from the cell body to the distal end of the axoneme, and IFT-A, which governs the retrograde return [13,55]. Even partial loss of IFT-B or IFT-A generally leads to impaired function of the entire subcomplex and loss of anterograde or retrograde functionality, respectively [56,57]. IFT has been shown to control cilia biogenesis and function in most ciliated eukaryotes, including vertebrates [58].

The CPLANE complex and IFT have been related, mainly with IFT-A. In the absence of the CPLANE complex, peripheral IFT-A proteins do not localise to basal bodies and do not assemble into the core of the IFT-A subcomplex [24] (Figure 1). Moreover, it has been observed that CPLANE does not move along axonemes, thus showing that it is not a component of the IFT particle itself. This fact seems logical, since very strong relationships have been observed between the proteins that form the complex, but a weak relationship has been observed between CPLANE proteins and IFT proteins [24].

Focusing on the different components of the CPLANE complex, the first indication that CPLANE proteins play an important role in IFT came from the observation that *FUZ* disruption specifically affected protein localisation in the distal, but not the proximal, axoneme in *Xenopus* multiciliated cells [20]. In another study, an effect of *FUZ* on intraflagellar transport was observed. Fuz protein is required for proper localisation of retrograde IFT proteins within the cytoplasm, thus allowing maintenance of the cilium [59].

As discussed above, Fuz interacts directly with Rsg1 [28], so an effect of Rsg1 on intraflagellar transport seems likely. Knockdown (KD) of *RSG1* function has been shown to lead to similar, but not identical, defects in axonemal IFT dynamics compared to the loss of *FUZ*. KD of *RSG1* has also been shown to lead to defects in cytoplasmic IFT organisation which are similar to those observed following *FUZ* disruption, and to disorganisation of

apically located basal bodies. However, the latter phenotype is not observed under *FUZ* KD conditions [27].




Figure 1. Relationship between the CPLANE complex and IFT. (A) Normal operation of the IFT when the CPLANE complex is undamaged; (B) IFT is truncated when the CPLANE complex is inoperative, generating shorter cilia.

For other components of the CPLANE complex, another study confirmed the effect of *JBTS17* on intraflagellar transport. *JBTS17* knockdown specifically disrupted the recruitment of IFT-A subunits from the peripheral zone to the basal body. The levels of Ift139, Ift121 and Ift43 in the basal bodies were dramatically reduced after *JBTS17* knockdown [24].

Regarding the effect of the CPLANE complex on the IFT-B subunit, it has been observed that disturbance of either *JBTS17* or *WDPCP* disrupted IFT-B movement [24], as it happens with *FUZ* [59].

These data argue that CPLANE facilitates IFT-A recruitment and assembly at basal bodies [24]. However, it is surprising that the role of *INTU* in intraflagellar transport is not known, just as *WDPCP*, which information in this regard is scarce. In this way, considering the clear effect of the rest of the components of the CPLANE complex on intraflagellar transport, it would be interesting to carry out future research on the effect of *INTU* and *WDPCP* on IFT.

5.2. Planar Cell Polarity

The planar cell polarity (PCP) cascade is a conserved signalling pathway that governs the polarisation of cells within the plane of a cell sheet [21].

PCP genes have been extensively studied in *Drosophila melanogaster*. In this regard, *Drosophila* PCP genes are grouped into core PCP genes and tissue-specific PCP effector genes. The genes that are encompassed in central PCP are *FRIZZLED* (*fz*), *DISHEVELLED* (*dsh*), *PRICKLE* (*pk*), *DIEGO* (*dgo*), *STRABISMUS* (*stbm*, or Van Gogh (*Vang*)) and *FLAMINGO* (*fmi*, or starry night (*stan*)). PCP effector genes include *INTURNED* (*in*), *FUZZY* (*fy*), *FRITZ* (*WDPCP*, *frtz*), and *NEMO* (*nmo*) [60–62]. In vertebrates, PCP genes (*INTURNED*, *FUZZY* and *WDPCP*) are involved in a variety of functions, including neural tube closure, inner ear sensory cell hair bundle orientation and hair follicle orientation [63]. It has been shown that disruption of core PCP genes often results in severe defects in these processes [63–65], whereas mutations in PCP effector genes generally result in more subtle defects [21,66]. It is therefore proposed that, unlike PCP core genes, PCP effector genes may act in a more localised manner [67]. In addition, some vertebrate animals, like mammals, have acquired more sophisticated developmental processes, as more complex organogenesis and have even more PCP genes than *Drosophila* [63], supporting the theory that different

PCP components are involved in different developmental processes in a tissue- or organ-specific manner.

The effect of genes belonging to the CPLANE complex on PCP has been studied. Disruption of *FUZ*, a PCP effector gene, resulted in delayed and arrested hair follicle development in mice, a phenotype not previously associated with PCP genes [67].

In another study carried out on *Xenopus laevis*, *INTURNED* and *FUZZY* were found to affect PCP signalling and cell intercalation [21], a fact that was expected because of their influence on planar cell polarity in the fly [22,66]. In particular, the directed elongation of microtubules is a critical aspect of both PCP in *D. melanogaster* and ciliogenesis [68]; because of this, it has been suggested that these proteins may influence the actin cytoskeleton during PCP in organising microtubules [21].

Finally, *Wdpcp* has been shown to modulate the actin cytoskeleton in mice by mediating the interaction of *Sept2*, a protein involved in the formation of cytoskeletal filaments, with actin. The formation of a *Wdpcp-Sept2* complex may be required for *Sept2* binding to and stabilisation of actin filaments. These results suggest that *WDPCP* may regulate planar cell polarity by modulating both the microfilament and microtubule cytoskeleton [33].

5.3. Hedgehog Signalling Pathway

The Hedgehog (Hh) signalling pathway is known to play an important role in embryonic development, organogenesis and tissue homeostasis [69]. In the adult stage, its activity is downregulated in most organs but can be reactivated in physiological and pathological processes such as tissue regeneration and cancer [70].

Numerous studies have long demonstrated the necessity of cilia for proper reception of the Hedgehog signal [71–74]. Activation of the Hh signalling pathway depends on the primary cilium, which is necessary for the translocation of Hh pathway components and the subsequent activation of Hh target genes, such as *GLI1* and *PTCH1* [75,76]. The importance of proper functioning of primary cilia for the Hh signalling pathway is well established for processes such as morphogenesis [3,77] or Hh-dependent tumour formation [78].

The effect of the CPLANE complex genes on the Hh signalling pathway has been extensively studied. It has been observed that up-regulation of *INTU* is necessary, but not sufficient, for ciliogenesis and oncogenic Hh signalling during basal cell carcinoma (BCC) formation [78]. *INTU* homozygous mutant mouse embryos show multiple defects, probably due to defective Hh signalling and *GLI3* processing [31]. In another study on mouse skin, suppression of Hh pathway activation was observed in *INTU* null mutant cells [32].

INTURNED and *FUZZY* function is required in the response to Hedgehog signals during early development in mammals [21]. In contrast, cilia are not required for Hedgehog signalling in *D. melanogaster* [79].

FUZ mutants in mice, generated by a gene-trap approach, exhibit multiple defects, including spinal cord and limb defects. In addition, these mutants have impaired Hh signalling and proteolytic processing of *GLI3*, which is consistent with the critical role of the Hh signalling pathway in the spinal cord and limb development. In this study, fewer cilia were observed in *FUZ* mutant mice, consistent with the key role of cilia in mediating Hh signalling in mammals [30]. Another study demonstrated the importance of *FUZ* in the formation of primary cilia in epidermal keratinocytes and dermal fibroblasts, both of which are required for Hh signalling during hair follicle morphogenesis [67].

Generation of a knockout mouse for *WDPCP* established that the CPLANE *Wdpcp* protein is essential for the development of the mouse appendicular skeleton, both in the limb bud and at later stages of skeletal development, including growth plate organisation. Significant alteration in hedgehog signalling activation was shown to be associated with defective proliferation and differentiation of skeletal tissues [80]. In *WDPCP* mutant MEFs was observed that *Wdpcp* deficiency disrupted *Shh* signalling, but in contrast, loss of *Wdpcp* function partially rescued the severe defect phenotypes of the *PTCH1* or *SMO*

knockout embryos, which seems to indicate that *Wdpcp* may constrain *Shh* signalling downstream of *Smo/Ptch1* [33].

The effect of *RSG1* on the Hedgehog pathway is not entirely clear. A recent study showed that the phenotypes of *RSG1* mutant mice could be caused by disruption of cilia-dependent Hh signalling. *GLI1*, a target of SHH signalling, was observed to be expressed at lower levels in the mutants, suggesting that *RSG1* acts downstream of SHH and upstream of *GLI1* in the central Hh signalling pathway [19]. As discussed above, *RSG1* mutants produce fewer hair cells, but these are of normal length. However, unlike the gene expression in WT MEFs, *GLI1* and *PTCH1* mRNA were not up-regulated in response to SAG in *RSG1* mutants. This suggests that although in the small number of mutant hair cells that form, Hh pathway activation is normal, this is not sufficient for normal Hh pathway activation in the cell population as a whole [19].

6. Conclusions

The primary cilium is a structure that can be found in most vertebrate cells and incorrect formation of the cilium can lead to ciliopathy. Several processes and signalling pathways are involved during ciliogenesis that are essential for proper function. Recent studies have confirmed a key role of the CPLANE complex in these processes. Mutations in the genes that conform the complex have been found to disturb the processes required during ciliogenesis, and the clinical features are related to ciliopathies. Given these data, further studies on the CPLANE complex are required, as it could play a key role in the development of certain ciliopathies.

Author Contributions: J.E.M.-S. and D.V. designed the study and wrote the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by projects PI15/00049 and PI19/00332, funded by the Instituto de Salud Carlos III and co-funded by the European Union (ERDF/ESF, “A way to make Europe”/ “Investing in your future”). Xunta de Galicia provided additional support through the Centro de Investigación de Galicia (CINBIO 2019–2022; Ref. ED431G-2019/06) and the program Consolidación e estructuración de unidades de investigación competitivas (ED431C-2018/54).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Badano, J.L.; Mitsuma, N.; Beales, P.L.; Katsanis, N. The Ciliopathies: An Emerging Class of Human Genetic Disorders. *Annu. Rev. Genom. Hum. Genet.* **2006**, *7*, 125–148. [[CrossRef](#)] [[PubMed](#)]
2. Pazour, G.J.; Agrin, N.; Leszyk, J.; Witman, G.B. Proteomic analysis of a eukaryotic cilium. *J. Cell Biol.* **2005**, *170*, 103–113. [[CrossRef](#)] [[PubMed](#)]
3. Eggenschwiler, J.T.; Anderson, K.V. Cilia and Developmental Signaling. *Annu. Rev. Cell Dev. Biol.* **2007**, *23*, 345–373. [[CrossRef](#)] [[PubMed](#)]
4. Fry, A.M.; Leaper, M.J.; Bayliss, R. The primary cilium: Guardian of organ development and homeostasis. *Organogenesis* **2014**, *10*, 62–68. [[CrossRef](#)]
5. Wallmeier, J.; Nielsen, K.G.; Kuehni, C.E.; Lucas, J.S.; Leigh, M.W.; Zariwala, M.A.; Omran, H. Motile ciliopathies. *Nat. Rev. Dis. Primer* **2020**, *6*, 77. [[CrossRef](#)]
6. Nishimura, Y.; Yamakawa, D.; Uchida, K.; Shiromizu, T.; Watanabe, M.; Inagaki, M. Primary cilia and lipid raft dynamics. *Open Biol.* **2021**, *11*, 210130. [[CrossRef](#)]
7. Scherft, J.P.; Daems, W.T. Single cilia in chondrocytes. *J. Ultrastruct. Res.* **1967**, *19*, 546–555. [[CrossRef](#)]
8. Rich, D.R.; Clark, A.L. Chondrocyte primary cilia shorten in response to osmotic challenge and are sites for endocytosis. *Osteoarthr. Cartil.* **2012**, *20*, 923–930. [[CrossRef](#)]
9. Berbari, N.F.; O’Connor, A.K.; Haycraft, C.J.; Yoder, B.K. The Primary Cilium as a Complex Signaling Center. *Curr. Biol.* **2009**, *19*, R526–R535. [[CrossRef](#)]

10. Izawa, I.; Goto, H.; Kasahara, K.; Inagaki, M. Current topics of functional links between primary cilia and cell cycle. *Cilia* **2015**, *4*, 12. [[CrossRef](#)]
11. Wheway, G.; Nazlamova, L.; Hancock, J.T. Signaling through the Primary Cilium. *Front. Cell Dev. Biol.* **2018**, *6*, 8. [[CrossRef](#)]
12. Nachury, M.V.; Seeley, E.S.; Jin, H. Trafficking to the Ciliary Membrane: How to Get Across the Periciliary Diffusion Barrier? *Annu. Rev. Cell Dev. Biol.* **2010**, *26*, 59–87. [[CrossRef](#)]
13. Ishikawa, H.; Marshall, W.F. Ciliogenesis: Building the cell's antenna. *Nat. Rev. Mol. Cell Biol.* **2011**, *12*, 222–234. [[CrossRef](#)]
14. Chen, H.Y.; Kelley, R.A.; Li, T.; Swaroop, A. Primary cilia biogenesis and associated retinal ciliopathies. *Semin. Cell Dev. Biol.* **2021**, *110*, 70–88. [[CrossRef](#)]
15. Reiter, J.F.; Leroux, M.R. Genes and molecular pathways underpinning ciliopathies. *Nat. Rev. Mol. Cell Biol.* **2017**, *18*, 533–547. [[CrossRef](#)]
16. Butler, M.T.; Wallingford, J.B. Planar cell polarity in development and disease. *Nat. Rev. Mol. Cell Biol.* **2017**, *18*, 375–388. [[CrossRef](#)]
17. Skoda, A.M.; Simovic, D.; Karin, V.; Kardum, V.; Vranic, S.; Serman, L. The role of the Hedgehog signaling pathway in cancer: A comprehensive review. *Bosn. J. Basic Med. Sci.* **2018**, *18*, 8–20. [[CrossRef](#)]
18. Varjosalo, M.; Taipale, J. Hedgehog: Functions and mechanisms. *Genes Dev.* **2008**, *22*, 2454–2472. [[CrossRef](#)]
19. Agbu, S.O.; Liang, Y.; Liu, A.; Anderson, K.V. The small GTPase RSG1 controls a final step in primary cilia initiation. *J. Cell Biol.* **2018**, *217*, 413–427. [[CrossRef](#)]
20. Gray, R.S.; Abitua, P.B.; Wlodarczyk, B.J.; Szabo-Rogers, H.L.; Blanchard, O.; Lee, I.; Weiss, G.S.; Liu, K.J.; Marcotte, E.M.; Wallingford, J.B.; et al. The planar cell polarity effector Fuz is essential for targeted membrane trafficking, ciliogenesis and mouse embryonic development. *Nat. Cell Biol.* **2009**, *11*, 1225–1232. [[CrossRef](#)]
21. Park, T.J.; Haigo, S.L.; Wallingford, J.B. Ciliogenesis defects in embryos lacking inturned or fuzzy function are associated with failure of planar cell polarity and Hedgehog signaling. *Nat. Genet.* **2006**, *38*, 303–311. [[CrossRef](#)]
22. Adler, P.N.; Lee, H. Frizzled signaling and cell–cell interactions in planar polarity. *Curr. Opin. Cell Biol.* **2001**, *13*, 635–640. [[CrossRef](#)]
23. Kim, S.K.; Shindo, A.; Park, T.J.; Oh, E.C.; Ghosh, S.; Gray, R.S.; Lewis, R.A.; Johnson, C.A.; Attie-Bittach, T.; Katsanis, N.; et al. Planar Cell Polarity Acts Through Septins to Control Collective Cell Movement and Ciliogenesis. *Science* **2010**, *329*, 1337–1340. [[CrossRef](#)]
24. Toriyama, M.; Lee, C.; Taylor, S.P.; Duran, I.; Cohn, D.H.; Bruel, A.-L.; Tabler, J.M.; Drew, K.; Kelly, M.R.; Kim, S.; et al. The ciliopathy-associated CPLANE proteins direct basal body recruitment of intraflagellar transport machinery. *Nat. Genet.* **2016**, *48*, 648–656. [[CrossRef](#)]
25. Lopez, E.; Thauvin-Robinet, C.; Reversade, B.; Khartoufi, N.E.; Devisme, L.; Holder, M.; Ansart-Franquet, H.; Avila, M.; Lacombe, D.; Kleinfinger, P.; et al. C5orf42 is the major gene responsible for OFD syndrome type VI. *Hum. Genet.* **2014**, *133*, 367–377. [[CrossRef](#)]
26. Alazami, A.M.; Alshammari, M.J.; Salih, M.A.; Alzahrani, F.; Hijazi, H.; Seidahmed, M.Z.; Abu Safieh, L.; Aldosary, M.; Khan, A.O.; Alkuraya, F.S. Molecular characterization of Joubert syndrome in Saudi Arabia. *Hum. Mutat.* **2012**, *33*, 1423–1428. [[CrossRef](#)]
27. Brooks, E.R.; Wallingford, J.B. The Small GTPase Rsg1 is important for the cytoplasmic localization and axonemal dynamics of intraflagellar transport proteins. *Cilia* **2013**, *2*, 13. [[CrossRef](#)] [[PubMed](#)]
28. Langousis, G.; Cavadini, S.; Boegholm, N.; Lorentzen, E.; Kempf, G.; Matthias, P. Structure of the ciliogenesis-associated CPLANE complex. *Sci. Adv.* **2022**, *8*, eabn0832. [[CrossRef](#)]
29. Gerondopoulos, A.; Strutt, H.; Stevenson, N.L.; Sobajima, T.; Levine, T.P.; Stephens, D.J.; Strutt, D.; Barr, F.A. Planar Cell Polarity Effector Proteins Inturned and Fuzzy Form a Rab23 GEF Complex. *Curr. Biol.* **2019**, *29*, 3323–3330.e8. [[CrossRef](#)]
30. Heydeck, W.; Zeng, H.; Liu, A. Planar cell polarity effector gene *Fuzzy* regulates cilia formation and Hedgehog signal transduction in mouse. *Dev. Dyn.* **2009**, *238*, 3035–3042. [[CrossRef](#)]
31. Zeng, H.; Hoover, A.N.; Liu, A. PCP effector gene Inturned is an important regulator of cilia formation and embryonic development in mammals. *Dev. Biol.* **2010**, *339*, 418–428. [[CrossRef](#)] [[PubMed](#)]
32. Dai, D.; Li, L.; Huebner, A.; Zeng, H.; Guevara, E.; Claypool, D.J.; Liu, A.; Chen, J. Planar cell polarity effector gene *Intu* regulates cell fate-specific differentiation of keratinocytes through the primary cilia. *Cell Death Differ.* **2013**, *20*, 130–138. [[CrossRef](#)] [[PubMed](#)]
33. Cui, C.; Chatterjee, B.; Lozito, T.P.; Zhang, Z.; Francis, R.J.; Yagi, H.; Swanhart, L.M.; Sanker, S.; Francis, D.; Yu, Q.; et al. Wdpcp, a PCP Protein Required for Ciliogenesis, Regulates Directional Cell Migration and Cell Polarity by Direct Modulation of the Actin Cytoskeleton. *PLoS Biol.* **2013**, *11*, e1001720. [[CrossRef](#)] [[PubMed](#)]
34. Waters, A.M.; Beales, P.L. Ciliopathies: An expanding disease spectrum. *Pediatr. Nephrol.* **2011**, *26*, 1039–1056. [[CrossRef](#)]
35. Quinlan, R.J.; Tobin, J.L.; Beales, P.L. Chapter 5 Modeling Ciliopathies. In *Mouse Models of Developmental Genetic Disease. Current Topics in Developmental Biology*; Elsevier: Amsterdam, The Netherlands, 2008; Volume 84, pp. 249–310. ISBN 978-0-12-374454-8.
36. Seo, J.H.; Zilber, Y.; Babayeva, S.; Liu, J.; Kyriakopoulos, P.; De Marco, P.; Merello, E.; Capra, V.; Gros, P.; Torban, E. Mutations in the planar cell polarity gene, *Fuzzy*, are associated with neural tube defects in humans. *Hum. Mol. Genet.* **2011**, *20*, 4324–4333. [[CrossRef](#)]

37. Zhang, W.; Taylor, S.P.; Ennis, H.A.; Forlenza, K.N.; Duran, I.; Li, B.; Sanchez, J.A.O.; Nevarez, L.; Nickerson, D.A.; Bamshad, M.; et al. Expanding the genetic architecture and phenotypic spectrum in the skeletal ciliopathies. *Hum. Mutat.* **2018**, *39*, 152–166. [[CrossRef](#)]
38. Huber, C.; Cormier-Daire, V. Ciliary disorder of the skeleton. *Am. J. Med. Genet. C Semin. Med. Genet.* **2012**, *160C*, 165–174. [[CrossRef](#)]
39. Wang, I.-Y.; Chung, C.-F.; Babayeva, S.; Sogomonian, T.; Torban, E. Loss of Planar Cell Polarity Effector Fuzzy Causes Renal Hypoplasia by Disrupting Several Signaling Pathways. *J. Dev. Biol.* **2021**, *10*, 1. [[CrossRef](#)]
40. Bruel, A.-L.; Franco, B.; Duffourd, Y.; Thevenon, J.; Jago, L.; Lopez, E.; Deleuze, J.-F.; Doummar, D.; Giles, R.H.; Johnson, C.A.; et al. Fifteen years of research on oral–facial–digital syndromes: From 1 to 16 causal genes. *J. Med. Genet.* **2017**, *54*, 371–380. [[CrossRef](#)]
41. Bruel, A.-L.; Levy, J.; Elenga, N.; Defo, A.; Favre, A.; Lucron, H.; Capri, Y.; Perrin, L.; Passemard, S.; Vial, Y.; et al. INTU-related oral-facial-digital syndrome type VI: A confirmatory report. *Clin. Genet.* **2018**, *93*, 1205–1209. [[CrossRef](#)]
42. Yakar, O.; Tatar, A. INTU-related oral-facial-digital syndrome XVII: Clinical spectrum of a rare disorder. *Am. J. Med. Genet. A* **2022**, *188*, 590–594. [[CrossRef](#)]
43. Shaheen, R.; Faqeh, E.; Alshammari, M.J.; Swaid, A.; Al-Gazali, L.; Mardawi, E.; Ansari, S.; Sogaty, S.; Seidahmed, M.Z.; AlMotairi, M.I.; et al. Genomic analysis of Meckel–Gruber syndrome in Arabs reveals marked genetic heterogeneity and novel candidate genes. *Eur. J. Hum. Genet.* **2013**, *21*, 762–768. [[CrossRef](#)]
44. Shamseldin, H.E.; Shaheen, R.; Ewida, N.; Bubshait, D.K.; Alkuraya, H.; Almardawi, E.; Howaidi, A.; Sabr, Y.; Abdalla, E.M.; Alfaifi, A.Y.; et al. The morbid genome of ciliopathies: An update. *Genet. Med.* **2020**, *22*, 1051–1060. [[CrossRef](#)]
45. Khan, S.A.; Muhammad, N.; Khan, M.A.; Kamal, A.; Rehman, Z.U.; Khan, S. Genetics of human Bardet-Biedl syndrome, an updates: Genetics of human Bardet-Biedl syndrome. *Clin. Genet.* **2016**, *90*, 3–15. [[CrossRef](#)]
46. M’hamdi, O.; Ouertani, I.; Chaabouni-Bouhamed, H. Update on the Genetics of Bardet-Biedl Syndrome. *Mol. Syndromol.* **2014**, *5*, 51–56. [[CrossRef](#)]
47. Hartill, V.; Szymanska, K.; Sharif, S.M.; Wheway, G.; Johnson, C.A. Meckel–Gruber Syndrome: An Update on Diagnosis, Clinical Management, and Research Advances. *Front. Pediatr.* **2017**, *5*, 244. [[CrossRef](#)]
48. Romani, M.; Mancini, F.; Micalizzi, A.; Poretti, A.; Miccinilli, E.; Accorsi, P.; Avola, E.; Bertini, E.; Borgatti, R.; Romaniello, R.; et al. Oral-facial-digital syndrome type VI: Is C5orf42 really the major gene? *Hum. Genet.* **2015**, *134*, 123–126. [[CrossRef](#)]
49. Srouf, M.; Schwartzentruber, J.; Hamdan, F.F.; Ospina, L.H.; Patry, L.; Labuda, D.; Massicotte, C.; Dobrzniecka, S.; Capo-Chichi, J.-M.; Papillon-Cavanagh, S.; et al. Mutations in C5ORF42 Cause Joubert Syndrome in the French Canadian Population. *Am. J. Hum. Genet.* **2012**, *90*, 693–700. [[CrossRef](#)]
50. Bayram, Y.; Aydin, H.; Gambin, T.; Akdemir, Z.C.; Atik, M.M.; Karaca, E.; Karaman, A.; Pehlivan, D.; Jhangiani, S.N.; Gibbs, R.A.; et al. Exome sequencing identifies a homozygous C5orf42 variant in a Turkish kindred with oral-facial-digital syndrome type VI. *Am. J. Med. Genet. A* **2015**, *167*, 2132–2137. [[CrossRef](#)]
51. Bachmann-Gagescu, R.; Dempsey, J.C.; Phelps, I.G.; O’Roak, B.J.; Knutzen, D.M.; Rue, T.C.; Ishak, G.E.; Isabella, C.R.; Gorden, N.; Adkins, J.; et al. Joubert syndrome: A model for untangling recessive disorders with extreme genetic heterogeneity. *J. Med. Genet.* **2015**, *52*, 514–522. [[CrossRef](#)]
52. Liu, Q.; Wang, H.; Zhao, J.; Liu, Z.; Sun, D.; Yuan, A.; Luo, G.; Wei, W.; Hou, M. Four novel compound heterozygous mutations in C5orf42 gene in patients with pure and mild Joubert syndrome. *Int. J. Dev. Neurosci.* **2020**, *80*, 455–463. [[CrossRef](#)]
53. Mardani, R.; Taghizadeh, E.; Taheri, F.; Raeisi, M.; Karimzadeh, M.R.; Rostami, D.; Ferns, G.A.; Ghayour-Mobarhan, M. A novel variant in C5ORF42 gene is associated with Joubert syndrome. *Mol. Biol. Rep.* **2020**, *47*, 4099–4103. [[CrossRef](#)]
54. Pedersen, L.B.; Rosenbaum, J.L. Chapter Two Intraflagellar Transport (IFT). In *Ciliary Function in Mammalian Development. Current Topics in Developmental Biology*; Elsevier: Amsterdam, The Netherlands, 2008; Volume 85, pp. 23–61. ISBN 978-0-12-374453-1.
55. Iomini, C.; Babaev-Khaimov, V.; Sassaroli, M.; Piperno, G. Protein Particles in Chlamydomonas Flagella Undergo a Transport Cycle Consisting of Four Phases. *J. Cell Biol.* **2001**, *153*, 13–24. [[CrossRef](#)]
56. Follit, J.A.; Tuft, R.A.; Fogarty, K.E.; Pazour, G.J. The Intraflagellar Transport Protein IFT20 Is Associated with the Golgi Complex and Is Required for Cilia Assembly. *Mol. Biol. Cell* **2006**, *17*, 3781–3792. [[CrossRef](#)]
57. Qin, J.; Lin, Y.; Norman, R.X.; Ko, H.W.; Eggenschwiler, J.T. Intraflagellar transport protein 122 antagonizes Sonic Hedgehog signaling and controls ciliary localization of pathway components. *Proc. Natl. Acad. Sci. USA* **2011**, *108*, 1456–1461. [[CrossRef](#)]
58. Scholey, J.M.; Anderson, K.V. Intraflagellar Transport and Cilium-Based Signaling. *Cell* **2006**, *125*, 439–442. [[CrossRef](#)]
59. Brooks, E.R.; Wallingford, J.B. Control of vertebrate intraflagellar transport by the planar cell polarity effector Fuz. *J. Cell Biol.* **2012**, *198*, 37–45. [[CrossRef](#)]
60. Klein, T.J.; Mlodzik, M. Planar Cell Polarization: An Emerging Model Points in the Right Direction. *Annu. Rev. Cell Dev. Biol.* **2005**, *21*, 155–176. [[CrossRef](#)]
61. Zallen, J.A. Planar Polarity and Tissue Morphogenesis. *Cell* **2007**, *129*, 1051–1063. [[CrossRef](#)] [[PubMed](#)]
62. Strutt, D.; Warrington, S.J. Planar polarity genes in the *Drosophila* wing regulate the localisation of the FH3-domain protein Multiple Wing Hairs to control the site of hair production. *Development* **2008**, *135*, 3103–3111. [[CrossRef](#)] [[PubMed](#)]
63. Wang, Y.; Nathans, J. Tissue/planar cell polarity in vertebrates: New insights and new questions. *Development* **2007**, *134*, 647–658. [[CrossRef](#)] [[PubMed](#)]
64. Jones, C.; Chen, P. Planar cell polarity signaling in vertebrates. *BioEssays* **2007**, *29*, 120–132. [[CrossRef](#)] [[PubMed](#)]

65. Simons, M.; Mlodzik, M. Planar Cell Polarity Signaling: From Fly Development to Human Disease. *Annu. Rev. Genet.* **2008**, *42*, 517–540. [[CrossRef](#)] [[PubMed](#)]
66. Lee, H.; Adler, P.N. The Function of the *frizzled* Pathway in the Drosophila Wing Is Dependent on *inturned* and *fuzzy*. *Genetics* **2002**, *160*, 1535–1547. [[CrossRef](#)]
67. Dai, D.; Zhu, H.; Wlodarczyk, B.; Zhang, L.; Li, L.; Li, A.G.; Finnell, R.H.; Roop, D.R.; Chen, J. Fuz Controls the Morphogenesis and Differentiation of Hair Follicles through the Formation of Primary Cilia. *J. Investig. Dermatol.* **2011**, *131*, 302–310. [[CrossRef](#)]
68. Hagiwara, H.; Ohwada, N.; Takata, K. Cell Biology of Normal and Abnormal Ciliogenesis in the Ciliated Epithelium. In *International Review of Cytology*; Elsevier: Amsterdam, The Netherlands, 2004; Volume 234, pp. 101–141. ISBN 978-0-12-364638-5.
69. Barakat, M.T.; Humke, E.W.; Scott, M.P. Learning from Jekyll to control Hyde: Hedgehog signaling in development and cancer. *Trends Mol. Med.* **2010**, *16*, 337–348. [[CrossRef](#)]
70. Hui, C.; Angers, S. Gli Proteins in Development and Disease. *Annu. Rev. Cell Dev. Biol.* **2011**, *27*, 513–537. [[CrossRef](#)]
71. Huangfu, D.; Anderson, K.V. Cilia and Hedgehog Responsiveness in the Mouse. *Proc. Natl. Acad. Sci. USA* **2005**, *102*, 11325–11330. [[CrossRef](#)]
72. Huangfu, D.; Liu, A.; Rakeman, A.S.; Murcia, N.S.; Niswander, L.; Anderson, K.V. Hedgehog signalling in the mouse requires intraflagellar transport proteins. *Nature* **2003**, *426*, 83–87. [[CrossRef](#)]
73. Liu, A.; Wang, B.; Niswander, L.A. Mouse intraflagellar transport proteins regulate both the activator and repressor functions of Gli transcription factors. *Development* **2005**, *132*, 3103–3111. [[CrossRef](#)]
74. Corbit, K.C.; Aanstad, P.; Singla, V.; Norman, A.R.; Stainier, D.Y.R.; Reiter, J.F. Vertebrate Smoothed functions at the primary cilium. *Nature* **2005**, *437*, 1018–1021. [[CrossRef](#)] [[PubMed](#)]
75. Goetz, S.C.; Anderson, K.V. The primary cilium: A signalling centre during vertebrate development. *Nat. Rev. Genet.* **2010**, *11*, 331–344. [[CrossRef](#)] [[PubMed](#)]
76. Singla, V.; Reiter, J.F. The Primary Cilium as the Cell’s Antenna: Signaling at a Sensory Organelle. *Science* **2006**, *313*, 629–633. [[CrossRef](#)] [[PubMed](#)]
77. Oh, E.C.; Katsanis, N. Cilia in vertebrate development and disease. *Development* **2012**, *139*, 443–448. [[CrossRef](#)]
78. Yang, N.; Leung, E.L.-H.; Liu, C.; Li, L.; Eguether, T.; Jun Yao, X.-J.; Jones, E.C.; Norris, D.A.; Liu, A.; Clark, R.A.; et al. INTU is essential for oncogenic Hh signaling through regulating primary cilia formation in basal cell carcinoma. *Oncogene* **2017**, *36*, 4997–5005. [[CrossRef](#)]
79. Han, Y.-G.; Kwok, B.H.; Kernan, M.J. Intraflagellar Transport Is Required in Drosophila to Differentiate Sensory Cilia but Not Sperm. *Curr. Biol.* **2003**, *13*, 1679–1686. [[CrossRef](#)]
80. Langhans, M.T.; Gao, J.; Tang, Y.; Wang, B.; Alexander, P.; Tuan, R.S. Wdpcp regulates cellular proliferation and differentiation in the developing limb via hedgehog signaling. *BMC Dev. Biol.* **2021**, *21*, 10. [[CrossRef](#)]